SUBSTANCE USE DISORDER TOOL

DATE:					NAME:				
lcohol F10	Opioid F11	Cannabis F12	Sedative Hypnotic Anxiolytic F13	Cocaine F14	Stimulant F15	PCP F16	Tobacco/ Nicotine F17	Inhalant F18	Oth F19
	Consuming the substance in larger amounts and for a longer amount of time than intended. Persistent desire to cut down or regulate use. The individual may have unsuccessfully attempted to stop in the past. Spending a great deal of time obtaining, using, or recovering from the effects of substance use. Experiencing craving, a pressing desire to use the substance.								
	Substance use impairs ability to fulfill major obligations at work, school, or home. Continued use of the substance despite it causing significant social or interpersonal problems. Reduction or discontinuation of recreational, social, or occupational activities because of substance use.								
77510	USE	Persisten		use despite l	ally unsafe en knowledge tha			ate physical o	or
	PHARMACOLOGICAL CRITERIA	desired specific Withdra the subsidecume Note: Individual withdraw	effect, or th symptoms a wal: A collect tance decre nted withdra dividuals car	e usual dose at different r ction of signs ase. Individua awal sympton n have an SUE LO and 11) in t	creasingly hig has a reduced ates. and symptoms als are likely to ns from halluc with prescrip the context of	d effect; ind s that occurs seek the sul inogens, PCF tion medicat	when blood a bstance to relict or inhalants. tions, so tolera	ouild tolerand and tissue leve ieve symptom ance and	ce to
Mild: 2-3 criteria				Moderate: 4-5 criteria			Severe: 6-11 criteria		